

Duality of Hopf C^* -algebras

Chi-Keung Ng

Abstract

In this paper, we study the duality theory of Hopf C^* -algebras in a general “Hilbert-space-free” framework. Our particular interests are the “full duality” and the “reduced duality”. In order to study the reduced duality, we define the interesting notion of Fourier algebra of a general Hopf C^* -algebra. This study of reduced duality and Fourier algebra is found to be useful in the study of other aspects of Hopf C^* -algebras (see e.g. [12], [13] and [14]).

2000 AMS Mathematics Classification numbers: Primary 46L55; 46L89; Secondary 22D15

0 Introduction

The study of Hopf C^* -algebras began with group C^* -algebras. There is a strong relation between locally compact groups and commutative (or co-commutative) Hopf C^* -algebras (see [7] and [16]). The general (i.e. non-commutative and non-co-commutative) Hopf C^* -algebras are getting more attention with the different studies of “locally compact quantum groups” in the recent literatures (see e.g. [1], [2], [3], [5], [6], [8], [9], [17], [18] and [21]).

One important and interesting feature of Hopf C^* -algebras is their duality theory (which is also related to the interesting notion of “amenability” – see the last paragraph of this introduction). In the literatures, duality is usually tied up with the notion of *multiplicative unitaries*. The aim of this paper is to study the duality theory of Hopf C^* -algebras without referring explicitly to multiplicative unitary. The advantage of this approach is that there may not be a multiplicative unitary in the first place and different multiplicative unitaries may give “the same duality” between two Hopf C^* -algebras. Moreover, it is not clear the kind of conditions (e.g. regularity in [2] and manageability in [21]) required on those multiplicative unitaries (e.g. the tensor product of a regular multiplicative unitary with a manageable one may not be regular nor manageable any more).

In this article, we will give a general intrinsic framework for the duality of Hopf C^* -algebras (which includes all sort of interesting multiplicative unitaries) and will pay more attention to the “full duality” (or “universal duality”) and the “reduced duality”. The first one will be studied in Section 2 (where its existence and related topics will be discussed). To define the second duality, we will need some preparation. In particular, we need to have some understanding of the general duality framework, namely, the Fourier duality. Section 3 is devoted to the study of this.

The reduced duality will be studied in Section 4. In order to define this, a natural notion of the “Fourier algebra” of a Hopf C^* -algebra is required. This object is interesting as its definition is intrinsic (i.e. concerning just the structure of the given Hopf C^* -algebra without referring to any extra structure) but it is the “core” of the duality theory given by multiplicative unitaries. Using this, we obtained a kind of uniqueness result.

The study of duality theory in this paper is important for the understanding of “amenability” of general Hopf C^* -algebras. In fact, roughly speaking, we can think of amenable Hopf C^* -algebras as those whose “full duality” and “reduced duality” coincide. We gave a detail study of this in another paper ([12]). Moreover, the Fourier algebras and its relation with reduced duality (as given in Section 4) is also important for the study of amenability of Kac algebras and Kac systems (see [13] and [14]).

1 Preliminary and notations

We begin with the definition of Hopf C^* -algebras and Hopf von Neumann algebras.

Definition 1.1 *Let S be a C^* -algebra.*

- (a) *A non-degenerate $*$ -homomorphism from S to $M(S \otimes S)$ is said to be a comultiplication if $(\delta \otimes \text{id})\delta = (\text{id} \otimes \delta)\delta$. A $*$ -homomorphism $\varphi \in S^* \setminus \{0\}$ is called a coidentity for δ if $(\varphi \otimes \text{id})\delta = \text{id} = (\text{id} \otimes \varphi)\delta$.*
- (b) *S is called a pre-Hopf C^* -algebra if it has a comultiplication δ . Moreover, (S, δ) is said to be a Hopf C^* -algebra, if $\delta(S)(1 \otimes S) \subseteq S \otimes S$ and $\delta(S)(S \otimes 1) \subseteq S \otimes S$.*
- (c) *Let A be a C^* -algebra and S be a pre-Hopf C^* -algebra. Then a non-degenerate $*$ -homomorphism ϵ from A to $M(A \otimes S)$ is called a coaction if $(\epsilon \otimes \text{id}) \circ \epsilon = (\text{id} \otimes \delta) \circ \epsilon$ and $\epsilon(A)(1 \otimes S) \subseteq A \otimes S$.*

Similarly, a non-degenerate $*$ -homomorphism from S to $M(S \otimes_{\max} S)$ is called a *full comultiplication* if it satisfies the equality in Definition 1.1(a). S is said to be a *full Hopf C^* -algebra* if it has a comultiplication satisfying similar conditions as in Definition 1.1(b).

Remark 1.2 (a) *Similarly, we can define Hopf von Neumann algebras and their coactions.*

- (b) *Let A and B be C^* -algebras and let $\mathcal{A}$ and $\mathcal{B}$ be von Neumann algebras. Suppose that φ and ψ are $*$ -homomorphisms from A and B to $\mathcal{A}$ and $\mathcal{B}$ respectively such that they are non-degenerate (in the sense that the image of any approximate unit converges to 1 weakly). Then $\varphi \otimes \psi$ extended to a map on $M(A \otimes B)$. Note that for any $F \in (\overline{A \otimes B})_*$ and $m \in M(A \otimes B)$, we have $(\varphi \otimes \psi)^*(F \cdot (\varphi \otimes \psi)(m)) = (\varphi \otimes \psi)^*(F) \cdot m$.*
- (c) *By part (b) above, we see that if A is a C^* -algebra and S is a Hopf C^* -algebra, then S^{**} is a Hopf von Neumann algebra and any coaction ϵ on A by S induces a weakly continuous coaction ϵ^{**} on A^{**} by S^{**} .*

For any subspace $N \subseteq S^*$, we denote $N^\top = \{x \in S : f(x) = 0 \text{ for all } f \in N\}$.

Proposition 1.3 *Let (S, δ) be a Hopf C^* -algebra and e be a non-trivial central projection of S^{**} . Let $T = S/(e \cdot S^*)^\top$ with the canonical quotient map Q . Then the following statements are equivalent.*

- (1) $\ker(Q) \subseteq \ker(Q \otimes \text{id})\delta$;
- (2) δ induces a coaction ϵ on T by S (i.e. $\epsilon \circ Q = (Q \otimes \text{id}) \circ \delta$);
- (3) $\overline{e \cdot S^{*\sigma(S^*, S)}}$ is a right ideal of S^* .

Proof: It is clear that $\ker(Q) \subseteq \ker(Q \otimes \text{id})\delta$ implies that δ induces a coaction on T by S . If δ induces a coaction ϵ on T by S , then for any $f \in T^*$ and $g \in S^*$, $Q^*(f)g = (f \otimes g)(Q \otimes \text{id}) \circ \delta = Q^*(h)$ where $h = (f \otimes g) \circ \epsilon$. Hence $\overline{e \cdot S^{*\sigma(S^*, S)}} = Q^*(T^*)$ is a right ideal of S^* . Finally, assume that $\overline{e \cdot S^{*\sigma(S^*, S)}}$ is a right ideal of S^* . For any $x \in \ker(Q)$, we have $(Q^*(f)g)(x) = 0$ (for $f \in T^*$ and $g \in S^*$) and thus, $(Q \otimes \text{id})\delta(x) = 0$.

Remark 1.4 (a) There is a corresponding result for $\ker(Q) \subseteq \ker(\text{id} \otimes Q)\delta$.

(b) By a similar argument as that of Proposition 1.3, we can also show that the followings are equivalent:

- (1) $\ker(Q) \subseteq \ker(Q \otimes Q)\delta$;
- (2) δ induces a quotient Hopf C^* -algebra structure on T ;
- (3) $\overline{e \cdot S^{*\sigma(S^*, S)}}$ is a subalgebra of S^* .

(c) For any locally compact group G , the only quotient Hopf C^* -algebra of $C_r^*(G)$ that has a compatible coaction by $C_r^*(G)$ is $C_r^*(G)$ itself (combine part (b) and Proposition 1.3 with [4, 2.4]).

Proposition 1.5 Let $\mathcal{M}$ be a Hopf von Neumann algebra with a comultiplication $\bar{\delta}$ and $e \in \mathcal{M}$ be a non-trivial central projection. Let $\bar{Q}$ be the canonical map from $\mathcal{M}$ to $e\mathcal{M}$. Then the following statements are equivalent.

- (1) $\ker(\bar{Q}) \subseteq \ker(\bar{Q} \otimes \text{id})\bar{\delta}$;
- (2) $(\bar{Q} \otimes \text{id}) \circ \bar{\delta}|_{e\mathcal{M}}$ is a coaction on $e\mathcal{M}$ by $\mathcal{M}$;
- (3) $\bar{\delta}(e) \geq e \otimes 1$;
- (4) $e\mathcal{M}_*$ is a right ideal of $\mathcal{M}_*$.

Proof: Assume that (1) holds. Then there exists a weak- $*$ -continuous map $\bar{e}$ such that $\bar{e} \circ \bar{Q} = (\bar{Q} \otimes \text{id}) \circ \bar{\delta}$. It is clear that $\bar{e}(e) = \bar{e}(\bar{Q}(1)) = e \otimes 1$ and $\bar{e} = (\bar{Q} \otimes \text{id}) \circ \bar{\delta}|_{e\mathcal{M}}$ is a coaction. Hence (2) is established. It is easy to see that (2) implies (3) (as $(\bar{Q} \otimes \text{id}) \circ \bar{\delta}(e) = e \otimes 1$). Suppose that (3) is true (i.e. $(e \otimes 1)\bar{\delta}(1 - e) = 0$). For any $z \in \mathcal{M}$ and $\omega, \nu \in \mathcal{M}_*$, we have $(1 - e)(e\omega \cdot \nu)(z) = (\omega \otimes \nu)(e \otimes 1)\bar{\delta}((1 - e)z) = 0$ and thus, (4) holds. Finally, we need to show that (4) implies (1). Let $z \in \ker(\bar{Q})$ (i.e. $ez = 0$). For any $\omega \in e\mathcal{M}_*$ and $\nu \in \mathcal{M}_*$, one has $(\omega \otimes \nu)(\bar{Q} \otimes \text{id})\bar{\delta}(z) = (e\omega \cdot \nu)(z) = 0$. This completes the proof.

Remark 1.6 (a) The equivalence of (3) and (4) is, in fact, [4, 2.2] in the case of central projections.

(b) We also have the von Neumann algebra version of Remark 1.4(b). In particular, the map $\bar{\delta}_e$ from $e\mathcal{M}$ to $e\mathcal{M} \otimes e\mathcal{M}$ defined by $\bar{\delta}_e(x) = (e \otimes e)\bar{\delta}(x)$ is a comultiplication on $e\mathcal{M}$ if and only if $\bar{\delta}(e) \geq e \otimes e$. It is the case if and only if $e\mathcal{M}_*$ is a subalgebra of $\mathcal{M}_*$.

(c) Let T be a quotient C^* -algebra of a Hopf C^* -algebra S and Q be the canonical quotient map. Then $Q^*(T^*)$ is both S -invariant and $\sigma(S^*, S)$ -closed. If e_T is a central projection in S^{**} such that $Q^*(T^*) = e_T \cdot S^*$ (as $Q^*(T^*)$ is S -invariant), then Propositions 1.3 and 1.5 as well as Remark 1.4(b) and part (b) above give relations between the properties of T and the properties of e_T .

We recall the following notations from [20]: for any Hilbert space H and any C^* -algebra A , let $C^*(H)$ be the collection of all non-degenerate separable C^* -subalgebras of $\mathcal{L}(H)$ and $\text{Rep}(A, H)$ be the collection of all non-degenerate representations of A on H .

Definition 1.7 ([20, 4.1]) Let S and T be two separable C^* -algebras and $X\eta(T \otimes S)$ (please refer to [20] for its meaning). Then S is said to be generated by X if for any separable Hilbert space H , any $B \in C^*(H)$ and any $\pi \in \text{Rep}(S, H)$, condition (i) below implies condition (ii):

- (i) $(\text{id} \otimes \pi)(X)\eta(T \otimes B)$;
- (ii) $\pi(S)B$ is dense in B .

The following simple result can be found in the second paragraph after [20, 4.1].

Lemma 1.8 Let H be a separable Hilbert space and C be a separable C^* -algebra. If $X\eta(C \otimes \mathcal{K}(H))$, then there exists at most one $A \in C^*(H)$ such that $X\eta(C \otimes A)$ and A is generated by X .

Proposition 1.9 *Let A, B, C and D be separable C^* -algebras. Let $X \in \eta(C \otimes A)$ such that A is generated by X .*

(a) Suppose that there is a surjective $$ -homomorphism φ from A to D . Then D is generated by $(\text{id} \otimes \varphi)(X)$.*

(b) Suppose that ϕ is a non-degenerate $$ -homomorphism from A to $M(B)$ such that B is generated by $(\text{id} \otimes \phi)(X)$. Then $\phi(A) = B$.*

Proof: (a) Let H be a Hilbert space. Suppose that $E \in C^*(H)$ and $\pi \in \text{Rep}(D, H)$ such that $(\text{id} \otimes \pi)(\text{id} \otimes \varphi)(X) \in \eta(C \otimes E)$. Since $\pi \circ \varphi \in \text{Rep}(A, H)$, $\pi(D)E = \pi \circ \varphi(A)E$ is dense in E .

(b) Let $D = \phi(A) \subseteq M(B)$ and φ be the canonical map from A to D . By part (a), D is generated by $(\text{id} \otimes \varphi)(X)$. Now, consider π to be a faithful non-degenerate representation of B on a Hilbert space H . Then π induces a faithful non-degenerate representation $\tilde{\pi}$ of $M(B)$ and hence of D . Let $Y = (\text{id} \otimes \pi \circ \phi)(X) = (\text{id} \otimes \tilde{\pi} \circ \varphi)(X) \in \eta(C \otimes \mathcal{K}(H))$. Since both $\pi(B)$ and $\tilde{\pi}(D)$ are generated by Y , we have, by Lemma 1.8, $\pi(B) = \tilde{\pi}(D)$ and by the injectivity of $\tilde{\pi}$, $B = D$.

2 The universal duality

In this section, we will study the universal object corresponding to unitary corepresentations of a Hopf C^* -algebra (which is an analogue of the full group C^* -algebras). Throughout this section, (S, δ) is a Hopf C^* -algebra.

Definition 2.1 *Let B be a C^* -algebra and H be a Hilbert space. A unitary $v \in M(B \otimes S)$ is called a unitary corepresentation of S in B (respectively, unitary corepresentation of S on H when $B = \mathcal{K}(H)$) if $(\text{id} \otimes \delta)(v) = v_{12}v_{13}$. A unitary corepresentation v of S on H is said to be cyclic if the $\varphi_v(S^*)$ is a cyclic representation on H (where $\varphi_v(f) = (\text{id} \otimes f)(v)$ for $f \in S^*$).*

We can define a sort of *universal dual* of S as a pair (B, v) where v is a unitary corepresentation of S in B such that for any unitary corepresentation u of S in any C^* -algebra D , there exists a unique non-degenerate $*$ -homomorphism φ from B to $M(D)$ with $u = (\varphi \otimes \text{id})(v)$. However, this type of objects is too abstract to handle and the following “stronger version” seems more useful (see e.g. [2] or [10]).

Definition 2.2 *$(\hat{S}_p, V_S)$ is said to be a strong dual of S if V_S is a unitary corepresentation of S in $\hat{S}_p$ such that*

- (1) for any unitary corepresentation u of S on a Hilbert space H , there exists a non-degenerate representation φ_u of $\hat{S}_p$ on H such that $u = (\varphi_u \otimes \text{id})(V_S)$;*
- (2) the set $\{(\text{id} \otimes f)(V_S) : f \in S^*\} \cap \hat{S}_p$ is dense in $\hat{S}_p$.*

Remark 2.3 *(a) If the strong dual of S exists, it is unique up to isomorphism.*

(b) It is clear that $S^\sim = \{f \in S^ : (\text{id} \otimes f)(V_S) \in \hat{S}_p\}$ is a closed ideal of S^* .*

The next proposition characterizes the existence of strong duals (c.f. [11, 2.1]). It requires the following well known lemma.

Lemma 2.4 *Let B be an algebra. Then there exists an injection from the collection of equivalent classes of cyclic representations of B (recall that a representation ϕ of B on a Hilbert space H is cyclic if $\phi(B)\xi$ is dense in H for some $\xi \in H$) to the set of positive sesquilinear forms on B .*

Proposition 2.5 *The strong dual of S exists if and only if there exists an ideal N of S^* such that for any unitary corepresentation X of S on H , the following conditions are satisfied:*

- (i) $\hat{S}_X = \{(\text{id} \otimes f)(X) : f \in N\}$ is a non-degenerate C^* -subalgebra of $\mathcal{L}(H)$;
- (ii) $X \in M(\hat{S}_X \otimes S)$.

Proof: Suppose that the strong dual of S exists and let $N = S^\sim$ (Remark 2.3(b)). Then for any unitary corepresentation X of S on H , there exists a representation φ of $\hat{S}_p$ on H such that $X = (\varphi \otimes \text{id})(V_S)$. Thus, we have $\hat{S}_X = \varphi(\hat{S}_p)$ and $X \in M(\hat{S}_X \otimes S)$. Conversely, suppose that there exists $N \subseteq S^*$ satisfying the two conditions. By definition, cyclic unitary corepresentations of S induce cyclic representations of S^* . Therefore, Lemma 2.4 ensures that the collection of all equivalent classes of cyclic unitary corepresentations is a set. It is now valid to consider the direct sum U of all equivalent classes of cyclic unitary corepresentations of S . It is obvious that U is a unitary corepresentation. For any given cyclic unitary corepresentation X of S on H , there exists a representation φ of $\hat{S}_U$ on H such that $X = (\varphi \otimes \text{id})(U)$. On the other hand, for any unitary corepresentation Y of S on H , since N is an ideal of S^* , we see that $\{(\text{id} \otimes g)(Y) : g \in S^*\} \subseteq M(\hat{S}_Y)$. Hence Y is cyclic if and only if $\hat{S}_Y$ acts cyclicly on H . Since $\hat{S}_Y$ is a C^* -algebra, the canonical representation of $\hat{S}_Y$ decomposes into a direct sum of cyclic representations of $\hat{S}_Y$. It is clear that each of these direct summands corresponds to a cyclic unitary corepresentation of S . Thus, there exists a representation φ of $\hat{S}_U$ on H such that $Y = (\varphi \otimes \text{id})(U)$.

Proposition 2.6 *Suppose that the strong dual $\hat{S}_p$ of S exists. There exists a full comultiplication $\hat{\delta}_p : \hat{S}_p \rightarrow M(\hat{S}_p \otimes_{\max} \hat{S}_p)$ such that $(\hat{\delta}_p \otimes \text{id})(V_S) = (V_S)_{13}(V_S)_{23} \in M((\hat{S}_p \otimes_{\max} \hat{S}_p) \otimes S)$. Moreover, there exists a coidentity for this comultiplication. If, in addition, $S^\sim$ is S -invariant (see Remark 2.3(b)), then $\hat{S}_p$ is a full Hopf C^* -algebra.*

The proofs of the first two statements are easy and the proof of the last one is similar to that of Proposition 3.3 in the next section. Moreover, by a similar argument as that of Proposition 3.3, if the set $\{(g \otimes \text{id})(V_S) : g \in \hat{S}_p^*\} \cap S$ is dense in S , then the converse of the last statement in the Proposition 2.6 also holds.

Definition 2.7 *A Hopf C^* -algebra S is said to be*

- (a) *dualizable if the strong dual $(\hat{S}_p, V_S)$ exists and $S^\sim$ is S -invariant;*
- (b) *symmetrically dualizable if it is dualizable and the set $\{(g \otimes \text{id})(V_S) : g \in \hat{S}_p^*\} \cap S$ is dense in S .*

Example 2.8 (a) *If V is a C^* -multiplicative unitary on H ([11, 2.3(b)]), then the Hopf C^* -algebra $S_V = \{(\omega \otimes \text{id})(W) : \omega \in \mathcal{L}(H)_*\}$ is symmetrically dualizable. This includes those Hopf C^* -algebras studied in [1], [2] and [21]. Moreover, as the “locally compact quantum groups” studied in [8] and [9] are defined by manageable multiplicative unitaries, they are also symmetrically dualizable. In particular, $C_0(G)$ and $C_r^*(G)$ are symmetrically dualizable. On the other hand, $C^*(G)$ is also symmetrically dualizable and more generally, $(S_V)_p$ is dualizable if V is a regular multiplicative unitary (see [2]).*

(b) *Let S be a unital C^* -algebra with the trivial comultiplication $\delta(a) = 1 \otimes a$. Then S is dualizable with $\hat{S}_p = \mathbf{C}$. S is certainly not symmetrically dualizable unless $S = \mathbf{C}$.*

(c) *Suppose that S and T are two Hopf C^* -algebras with coidentities e_S and e_T respectively. If both S and T are dualizable (respectively, symmetrically dualizable), then so is $S \otimes T$. In fact, if Z is a unitary corepresentation of $S \otimes T$, then $X = (\text{id} \otimes \text{id} \otimes e_T)(Z)$ and $Y = (\text{id} \otimes e_S \otimes \text{id})(Z)$ are unitary corepresentations of S and T respectively such that $X_{12}Y_{13} = Y_{13}X_{12}$.*

Remark 2.9 Given a coaction ϵ of a Hopf C^* -algebra S on a C^* -algebra A , one can define covariant representations for ϵ . Moreover, similar to the definition of the strong dual, one can also define the full crossed product of ϵ (which may or may not exist). It can be shown that if S is dualizable, then the full crossed product of ϵ exists if and only if there exists a covariant representation for ϵ . However, as these are not relevant to the main concern of this paper, we will not give the details here.

3 Fourier duality and representations

In this section, we will study a general duality framework.

Definition 3.1 Let S and T be Hopf C^* -algebras and $X \in M(T \otimes S)$ be a unitary. Then (T, X, S) is called a Fourier duality if

$$(\text{id} \otimes \delta_S)(X) = X_{12}X_{13} \quad \text{and} \quad (\delta_T \otimes \text{id})(X) = X_{13}X_{23} \quad (1)$$

as well as

$$\overline{\{(\text{id} \otimes f)(X) : f \in S^*\} \cap T} = T \quad \text{and} \quad \overline{\{(g \otimes \text{id})(X) : g \in T^*\} \cap S} = S.$$

Example 3.2 (a) For any C^* -multiplicative unitary V ([11, 2.1]), we recall that the following algebras $\hat{S}_V = \overline{\{(\text{id} \otimes \omega)(W) : \omega \in \mathcal{L}(H)_*\}}$ and $S_V = \overline{\{(\omega \otimes \text{id})(W) : \omega \in \mathcal{L}(H)_*\}}$ are Hopf C^* -algebras. Furthermore, $(\hat{S}_V, V, S_V)$ is a Fourier duality. In particular, it is the case for regular ([2, 3.3]), semi-regular and balanced ([1, 3.1]), or manageable ([21, 1.2] and also [9]) multiplicative unitaries.

(b) If (T, X, S) and (T', X', S') are two Fourier dualities, then so is $(T \otimes T', (X \otimes X')^{\sigma_{23}}, S \otimes S')$ where σ_{23} is the flips of the two middle variables. In particular, if V and W are multiplicative unitaries such that V is regular and W is manageable, then $(\hat{S}_V \otimes \hat{S}_W, (V \otimes W)^{\sigma_{23}}, S_V \otimes S_W)$ is also a Fourier duality. However, there is no reason to believe that $(V \otimes W)^{\sigma_{23}}$ is either regular or manageable.

(c) If S is symmetrically dualizable (see Definition 2.7(b)), then $(\hat{S}_p, V_S, S)$ is a Fourier duality.

(d) Let V and W be two regular multiplicative unitaries. Then $((\hat{S}_V)_p \otimes \hat{S}_W, (V_{S_V} \otimes W)^{\sigma_{23}}, S_V \otimes S_W)$ is also a Fourier duality. In general, it is not possible to represent this sort of duality using multiplicative unitary unless V is amenable.

(e) Suppose that (T, X, S) is a Fourier duality. Then (S, X^σ, T) , $(T^{\text{op}}, X^*, S^{\text{op}})$ and $(S^{\text{op}}, X^\top, T^{\text{op}})$ are all Fourier dualities (where σ is the flip of variables, $X^\top = (X^\sigma)^*$ and S^{op} is the Hopf C^* -algebra which has the same underlying C^* -algebra as S but with the comultiplication $\delta_{S^{\text{op}}} = \sigma \circ \delta_S$). Note that in the setting of multiplicative unitaries, it is only valid to talk about X and $X^\top$ and the other two only appear when irreducibility is considered.

Suppose that S and T are pre-Hopf C^* -algebras (see Definition 1.1(b)) and $X \in M(T \otimes S)$ satisfying equalities (1). Let j_S and j_T be the canonical homomorphisms from T^* and S^* to $M(S)$ and $M(T)$ respectively i.e. $j_S(g) = (g \otimes \text{id})(X)$ and $j_T(f) = (\text{id} \otimes f)(X)$. We denote

$$S^\# = \{f \in S^* : j_T(f) \in T\} \quad \text{and} \quad T^\# = \{g \in T^* : j_S(g) \in S\}.$$

It is clear that $S^\#$ and $T^\#$ are closed ideals of S^* and T^* respectively.

Proposition 3.3 *Let T and S be pre-Hopf C^* -algebras. Suppose that $X \in M(T \otimes S)$ satisfies the conditions in Definition 3.1. Then $S^\#$ is S -invariant if and only if T is a Hopf C^* -algebra. Consequently, for any Fourier duality (T, X, S) , $S^\#$ and $T^\#$ are automatically S -invariant and T -invariant respectively.*

Proof: Suppose that T is a Hopf C^* -algebra. For any $f \in S^\#$ and $g \in T^\#$,

$$j_T(f \cdot j_S(g)) = (\text{id} \otimes f \cdot j_S(g))(X) = (g \otimes \text{id} \otimes f)(X_{13}X_{23}) = (g \otimes \text{id})(\delta_T(j_T(f)))$$

which is in T . Hence, $f \cdot j_S(g) \in S^\#$ and so $S^\# \cdot S \subseteq S^\#$ (by the density condition in Definition 3.1). Similarly, we have $j_T(j_S(g) \cdot f) = (\text{id} \otimes g)(\delta_T(j_T(f)))$ and $S \cdot S^\# \subseteq S^\#$. Conversely, suppose that $S^\#$ is S -invariant. For any $g' \in S^\#$, there exist $s \in S$ and $g \in S^\#$ such that $g' = s \cdot g$. Moreover, for any $t \in T$, $X(t \otimes s) \in T \otimes S$ can be approximated by elements of the form $\sum b_i \otimes a_i$ ($a_i \in S$; $b_i \in T$) and

$$\delta_T((\text{id} \otimes g')(X))(1 \otimes t) = (\text{id} \otimes \text{id} \otimes g)(X_{13}X_{23}(1 \otimes t \otimes s))$$

which can be approximated by $\sum (\text{id} \otimes b_i \cdot g)(X) \otimes a_i$. This shows that $\delta_T((\text{id} \otimes g')(X))(1 \otimes t) \in T \otimes T$ and thus, $\delta_T(T)(1 \otimes T) \subseteq T \otimes T$. On the other hand, by considering $g' = g \cdot s$, we see that $(t \otimes 1)\delta_T((\text{id} \otimes g')(X)) \in T \otimes T$ and so, $(T \otimes 1)\delta_T(T) \subseteq T \otimes T$.

Corollary 3.4 *If (T, X, S) is a Fourier duality, then $S^\#$ is invariant under the usual conjugation (which is given by $f^*(x) = f(x^*)$).*

Proof: Since $S^\#$ is S -invariant, by [15, III.4.4], any $f \in S^\#$ can be approximated by elements of the form $\sum_k s_k \cdot g_k$ where $s_k \in S$ and $g_k \in S^\# \cap S_+^*$. Therefore, f^* can be approximated by $\sum_k g_k \cdot s_k^* \in S^\#$.

If, in addition, S is dualizable, then we have two “dualities”: $(\hat{S}_p, V_S, S)$ and (T, X, S) . It is natural to ask whether there is any relation between T and $\hat{S}_p$. By definition, there exists a $*$ -homomorphism q_T from $\hat{S}_p$ to $M(T)$ such that $X = (q_T \otimes \text{id})(V_S)$. It is not hard to show that q_T is a Hopf $*$ -homomorphism. Moreover, the following result shows that the image of q_T is exactly T (which is a bit surprising in this general situation of Fourier duality).

Proposition 3.5 *Let (T, X, S) be a Fourier duality. If S is dualizable and q_T is the induced map from $\hat{S}_p$ to $M(T)$, then S is symmetrically dualizable (i.e. $(\hat{S}_p, V_S, S)$ is a Fourier duality) and $T = q_T(\hat{S}_p)$.*

Proof: S is symmetrically dualizable because $\{(h \otimes \text{id})(V_S) : h \in q_T^*(T^\#)\}$ is dense in S . Now, suppose that $A = q_T(\hat{S}_p)$. Since both $S^\sim$ (see Remark 2.3(b)) and $S^\#$ are ideals of S^* , $S^\# \cdot S^\sim \subseteq S^\sim \cap S^\#$. Therefore, $T \cdot A \subseteq A \cap T$. Moreover, since q_T is non-degenerate, $T \cdot A = T$. This implies that $T \subseteq A$. On the other hand, it is clear that $X = (q_T \otimes \text{id})(V_S) \in M(A \otimes S)$. Take any $h \in S^\#$ and any $s \in S$ such that $h(s) = 1$. For any $a \in A$, because $X(A \otimes S) = A \otimes S$, we can approximate $a \otimes s$ by elements of the form

$$\sum_i X((\text{id} \otimes f_i)(X) \otimes s_i) = \sum_i (\text{id} \otimes \text{id} \otimes f_i)(X_{12}X_{13})(1 \otimes s_i)$$

(where $f_i \in S^\sim$ and $s_i \in S$). Hence, a can be approximated by elements of the form

$$\sum_i (\text{id} \otimes s_i \cdot h)(X)(\text{id} \otimes f_i)(X) \in T \cdot A = T.$$

The next result gives a relation between Fourier duality and C^* -algebra generated by affiliated element. It is a generalization of [21, 1.6.6].

Proposition 3.6 *Let (T, X, S) be a Fourier duality such that both S and T are separable. Then S is generated by X in the sense of Woronowicz (see Definition 1.7) and T is generated by X^σ .*

Proof: Let H be a Hilbert space. Let $\varphi_1, \varphi_2 \in \text{Rep}(S, H)$ such that $(\text{id} \otimes \varphi_1)(X) = (\text{id} \otimes \varphi_2)(X)$. Since $\{f \in T^\# : (f \otimes \text{id})(X)\}$ is dense in S , we have $\varphi_1 = \varphi_2$. This shows that X satisfies condition I of [20, 4.2]. Next, suppose that $B \in C^*(H)$ and $\pi \in \text{Rep}(S, H)$ such that $(\text{id} \otimes \pi)(X) \eta(T \otimes B)$. Since X is bounded, $(\text{id} \otimes \pi)(X) \in M(T \otimes B)$. Moreover, as X is a unitary, $(\text{id} \otimes \pi)(X)(T \otimes B)$ is dense in $T \otimes B$. We now take any $\omega \in T^\# \setminus (0)$. As $T^\#$ is T -invariant, $T \cdot \omega \subseteq T^\#$ and hence $(\omega \otimes \text{id})(X(t \otimes 1)) = (t \cdot \omega \otimes \text{id})(X) \in S$. By putting $F = T$ and $r = X$ in [20, 4.2(II)], we see that S is generated by X . Similarly, T is generated by X^σ .

The argument in Proposition 3.6 also shows that if S is separable and dualizable, then $\hat{S}_p$ is generated by V_S^σ . Hence, if S and T are separable, we can also obtain Proposition 3.5 by Propositions 1.9(b) and 3.6.

Remark 3.7 *Suppose that (T, X, S) is a Fourier duality.*

(a) *It may not be true in general that $S^\#$ separates points of S but we can always “quotient out the annihilators of $S^\#$ ”. More precisely, if $I = \bigcap_{f \in S^\#} \ker f$, then it can be shown that $S_0 = S/I$ is a Hopf C^* -algebra and $(T, (\text{id} \otimes q)(X), S_0)$ is a Fourier duality (where q is the quotient map from S to S_0) with $q^*(S_0^\#) = S^\#$. The same can be done on T .*

(b) *It is clear that $j_T(S^\#)$ is a subalgebra of T . An interesting question is when it will be a $*$ -subalgebra. Let us first consider the (unbounded) “coinvolution” κ_S from $j_S(T^\#)$ to S given by*

$$\kappa_S(j_S(g)) = (g \otimes \text{id})(X^*)$$

($g \in T^\#$) (it is well defined because of Corollary 3.4). Suppose that $S^\#$ and $T^\#$ separate points of T and S respectively. Then S can be regarded as a subalgebra of $(S^\#)^$. Moreover, we have the following equivalent statements:*

(i) *$j_T(S^\#)$ is a $*$ -subalgebra of T .*

(ii) *κ_S can be extended to a isometric anti-isomorphism from S onto S .*

(iii) *κ_S can be extended to a weak- $*$ -continuous anti-isomorphism from $S^{\#*}$ onto itself.*

Again, as this is not our main concern, we will not give the proof here.

Next, we turn to representations of a Fourier duality (T, X, S) . It is natural to start with unitary corepresentations of S and T and look for compatibility conditions with X .

Definition 3.8 *Let (T, X, S) be a Fourier duality.*

(a) *If u and v are unitary corepresentations of T and S respectively in B , then (u, v) is said to be a unitary X -covariant representation if it satisfies the following form of commutation relation:*

$$u_{12}^\sigma X_{13} v_{23} = v_{23} u_{12}^\sigma.$$

(b) *If μ and ν are representations of S and T respectively on H , then (μ, ν) is said to be a X -covariant representation if $((\mu \otimes \text{id})(X^\sigma), (\nu \otimes \text{id})(X))$ is a unitary X -covariant representation. Moreover, (μ, ν) is said to be faithful if and only if both μ and ν are injective.*

Remark 3.9 (a) Note that if X is any unitary in $M(T \otimes S)$ such that there exist unitary corepresentations u and v of T and S respectively satisfying the equality in Definition 3.8(a), then [11, 1.5] implies that X satisfies equalities (1).

(b) A unitary X^σ -covariant representation (v, u) means that $v_{12}^\sigma X_{13}^\sigma u_{23} = u_{23} v_{12}^\sigma$ or equivalently, $v_{23} X_{13} u_{12}^\sigma = u_{12}^\sigma v_{23}$. In general, there is no relation between X -covariant representations and X^σ -covariant representations.

(c) Let u and v be unitary corepresentations of S and T respectively on a Hilbert space H . Then (u, v) is a unitary X -covariant (respectively, X^* -covariant) representation if and only if (v, u) is a unitary $X^\top$ -covariant (respectively, X^σ -covariant) representation.

Proposition 3.10 If (μ, ν) is a X -covariant representation, then $W = (\nu \otimes \mu)(X)$ is a multiplicative unitary. Moreover, both $\mu(S)$ and $\nu(T)$ are Hopf C^* -algebras (with coactions induced by W), both μ and ν are Hopf $*$ -homomorphisms and $W \in M(\nu(T) \otimes \mu(S))$ satisfies equalities (1).

In fact, since $(\mu \otimes \mu)\delta_S(s) = W(\mu(s) \otimes 1)W^*$, the map $\bar{\delta}_S$ on $\mu(S)$ defined by $\bar{\delta}_S(\mu(s)) = (\mu \otimes \mu)\delta_S(s)$ is a well defined comultiplication.

Hence, (μ, ν) can be regarded as a sort of multiplicative unitary realization of the abstract Fourier duality (T, X, S) . However, $\mu(S)$ and $\nu(T)$ may not be defined by W in the usual way i.e. it is possible that $\nu(T) \neq \hat{S}_W$ or $\mu(S) \neq S_W$ (see Example 3.2(a)).

On the other hand, there may not exist any faithful X -covariant representation (e.g. if $(C^*(G), X, C_0(G))$ is the strong dual of $S = C_0(G)$ where G is a non-amenable locally compact group).

4 Reduced duality and uniqueness

In this section, we will define reduced Fourier dualities and study their uniqueness. This is done by using a notion of Fourier algebra of an arbitrary Hopf C^* -algebra. For a “nice” multiplicative unitary V , we can recover S_V from the structure of $(S_V)_p$ through this construction without looking at V (this fact is important for later applications in [12], [13] and [14]).

For simplicity, we call a subspace N of S^* a *Hopf ideal* (respectively, *left Hopf ideal* or *right Hopf ideal*) if it is a non-trivial S -invariant closed 2-sided ideal (respectively, left ideal or right ideal) of S^* . For any Fourier duality (T, X, S) , the subspace $S^\#$ is a Hopf ideal. Note that we use the term “Hopf ideal” in a different way from the literatures on Hopf algebras (although the quotient of S with respect to the annihilators of an Hopf ideal is a Hopf C^* -algebra).

Definition 4.1 (a) The Fourier algebra, A_S , of a Hopf C^* -algebra S is the intersection of all Hopf ideals of S^* .

(b) A Hopf C^* -algebra S is said to be *irreducible* if any Hopf ideal of S^* separates points of S .

(c) A Fourier duality (T, X, S) is said to be *reduced* if $\{(\text{id} \otimes f)(X) : f \in A_S\}$ is dense in T and $\{(g \otimes \text{id})(X) : g \in A_T\}$ is dense in S .

Note that the definition of the Fourier algebras comes from the corresponding property of $A(G)$ (see [4, 2.5]). In the following, we will show that it is really a “right definition”.

Lemma 4.2 *Let S be a irreducible Hopf C^* -algebra. Then its comultiplication δ_S is injective.*

Proof: Let I be the kernel of δ_S . It is clear that $I^\perp = \{f \in S^* : f(I) = 0\}$ is a S -invariant closed subspace of S^* . Moreover, for any $f, g \in S^*$, we have $fg \in I^\perp$ and in particular, $I^\perp$ is an ideal of S^* . Hence, either $I^\perp = (0)$ (i.e. $I = S$ and is impossible as δ is a non-degenerate $*$ -homomorphism) or $I^\perp$ is $\sigma(S^*, S)$ -dense in S^* which implies that $I^\perp = S^*$ (i.e. $I = (0)$).

It is natural to ask whether we can get a sort of “irreducible quotient” of a general Hopf C^* -algebra. If S is a Hopf C^* -algebra such that $A_S \neq (0)$, we can take the quotient $S_{irr} = S/(A_S)^\perp$ with the canonical quotient map Q_S .

Proposition 4.3 *Let S and T be Hopf C^* -algebras such that $A_S \neq (0)$ and $A_T \neq (0)$. Then S_{irr} is an irreducible Hopf C^* -algebra with $A_{S_{irr}} = A_S$. Moreover, if (T, X, S) is a Fourier duality and $X_{irr} = (Q_T \otimes Q_S)(X)$, then $(T_{irr}, X_{irr}, S_{irr})$ is a reduced Fourier duality. Consequently, if both S and T are irreducible with non-zero Fourier algebras, then (T, X, S) is automatically reduced.*

Proof: By Remark 1.4(b) (and the fact that Fourier algebras are invariant and hence are defined by central projections), S_{irr} and T_{irr} are both Hopf C^* -algebras. Since $S_{irr}^* \cong \overline{A_S}^{\sigma(S^*, S)}$, we have $A_{S_{irr}} = A_S$. It is clear that Q_S and Q_T are Hopf $*$ -homomorphisms and X_{irr} satisfies equalities (1). It is required to show that $(T_{irr}, X_{irr}, S_{irr})$ is a reduced Fourier duality. Let

$$N = \{g \in T^* : (g \otimes f)(X) = 0 \text{ for any } f \in A_S\}.$$

Then N is a closed subspace of T^* . Moreover, for any $g \in N$, $f \in A_S$ and $h \in S^\#$,

$$((j_T(h) \cdot g) \otimes f)(X) = (g \otimes fh)(X) = 0$$

(since A_S is an ideal of S^*) and similarly $(g \cdot j_T(h) \otimes f)(X) = (g \otimes hf)(X) = 0$. This shows that N is T -invariant. On the other hand, for any $g \in N$, $l \in T^*$ and $f \in A_S$,

$$(gl \otimes f)(X) = (g \otimes j_S(l) \cdot f)(X) = 0$$

(as A_S is S -invariant and hence is $M(S)$ -invariant) and $(lg \otimes f)(X) = 0$. Thus, either $N = (0)$ or N is a Hopf ideal (or equivalently, $A_T \subseteq N$). Suppose that $A_T \subseteq N$. Then $(g' \otimes f')(X_{irr}) = 0$ for all $g' \in A_T = A_{T_{irr}}$ and $f' \in A_S = A_{S_{irr}}$. This implies that $X_{irr} = 0$ (as $A_{S_{irr}}$ and $A_{T_{irr}}$ separate points of S_{irr} and T_{irr} respectively and are invariant) and is impossible because X_{irr} is a unitary. Therefore, $N = (0)$ and $\{(\text{id} \otimes f)(X) : f \in A_S\}$ is norm dense in T . Similarly, $\{(g \otimes \text{id})(X) : g \in A_T\}$ is dense in S .

From the proof of the above proposition, we see that if we consider the duality between S^* and T^* defined by X , then the “polar” of an ideal of T^* is S -invariant and the “polar” of a closed T -invariant subspace is an ideal of S^* .

Example 4.4 (a) *For any locally compact group G , $C_r^*(G)$ is an irreducible Hopf C^* -algebra (by [4, 2.4]). In fact, we have $C_r^*(G) = C^*(G)_{irr}$.*

(b) *If $S = S_V$ (see Example 3.2(a)) for a regular irreducible multiplicative unitary V , then $(\hat{S}_p)_{irr} = \hat{S}_V$ (see Corollary 4.10(c) below). It is an interesting fact since we can obtain the “reduced dual object” $\hat{S}_V$ as well as $V \in M(\hat{S}_V \otimes S_V)$ without knowing a priori that S comes from a Kac system nor assuming any special representation of S .*

In the following, we will give some uniqueness type results for reduced Fourier dualities (Theorem 4.13 and Corollary 4.14(b)). We will also consider their relations with multiplicative unitaries (Proposition 4.12(b) and Corollary 4.14(a)).

Lemma 4.5 *Let (μ, ν) be a X -covariant representation and e_μ be the support projection of the extension $\tilde{\mu}$ of μ on S^{**} . Then $\delta_S^{**}(e_\mu) \geq e_\mu \otimes 1$.*

Proof: Since (μ, ν) is a X -covariant representation,

$$(\mu \otimes \text{id})\delta_S(x) = (\nu \otimes \text{id})(X)(\mu(x) \otimes 1)(\nu \otimes \text{id})(X)^*$$

($x \in S$). Note that by the weak density of S in S^{**} , the same equation holds for $x \in S^{**}$ (with μ replaced by $\tilde{\mu}$ and δ_S by δ_S^{**}). Therefore, $(\tilde{\mu} \otimes \text{id})\delta_S^{**}(e_\mu) = 1$ and so $(\tilde{\mu} \otimes \text{id})((e_\mu \otimes 1)\delta_S^{**}(e_\mu)) = (\tilde{\mu} \otimes \text{id})(e_\mu \otimes 1)$. Since $\tilde{\mu}$ is injective on $e_\mu S^{**}$, we have $(e_\mu \otimes 1)\delta_S^{**}(e_\mu) = e_\mu \otimes 1$.

Proposition 4.6 *Let (T, X, S) be a Fourier duality and p be a non-zero central projection in S^{**} . Let (μ, ν) be a X -covariant representation. If $\delta_S^{**}(p) \geq 1 \otimes p$, then $p \geq e_\mu$.*

Proof: Let $W = (\nu \otimes \mu)(X) \in \nu(T)'' \bar{\otimes} \mu(S)''$. Then, by Proposition 3.10, $(\tilde{\mu} \otimes \tilde{\mu})\delta_S^{**}(s) = W(\tilde{\mu}(s) \otimes 1)W^*$ for any $s \in S^{**}$ (where $\tilde{\mu}$ is as in the previous lemma). If $\tilde{p} = \tilde{\mu}(p)$, then $\tilde{p}$ is a central projection in $\mu(S)''$ and we have

$$\begin{aligned} W((1 - \tilde{p}) \otimes \tilde{p})W^* &= W(1 \otimes \tilde{p})W^* - (1 \otimes \tilde{p})W(\tilde{p} \otimes 1)W^* \\ &= 1 \otimes \tilde{p} - (\tilde{\mu} \otimes \tilde{\mu})((1 \otimes p)\delta_S^{**}(p)) = 0. \end{aligned}$$

Hence either $\tilde{p} = 0$ or $\tilde{p} = 1$. Suppose that $\tilde{\mu}(p) = \tilde{p} = 0$. Then $p \leq 1 - e_\mu$ and, by Lemma 4.5, $1 \otimes p + e_\mu \otimes 1 \leq \delta_S^{**}(p) + \delta_S^{**}(e_\mu) \leq 1 \otimes 1$. Therefore, $e_\mu \otimes 1 \leq 1 \otimes (1 - p)$ and so either $p = 0$ or $e_\mu = 0$ which contradicts the assumption. Thus, we have $\tilde{p} = 1$ or, equivalently, $p \geq e_\mu$.

Remark 4.7 (a) *The idea of the proof of Proposition 4.6 is mainly from that of [4, 2.4] but since we consider central projections only, we have a simpler proof here.*

(b) *We also have a corresponding result for ν in Lemma 4.5 and Proposition 4.6 (namely, $\delta_T^{**}(e_\nu) \geq 1 \otimes e_\nu$ and $\delta_T^{**}(q) \geq q \otimes 1$ implies that $q \geq e_\nu$).*

Corollary 4.8 *Let (T, X, S) be a Fourier duality such that $A_S \neq (0)$ and let (μ, ν) be a X -covariant representation. Then $e_\mu \cdot S^* \subseteq A_S$ and μ induces a representation μ_{irr} of S_{irr} such that (μ_{irr}, ν) is a $(\text{id} \otimes Q_S)(X)$ -covariant representation.*

Proof: Let p be the central projection of S^{**} such that $A_S = p \cdot S^*$ (as A_S is S -invariant). Then by Propositions 4.6 and 1.5 (note that A_S is a left ideal of S^*), $p \geq e_\mu$. Hence $e_\mu \cdot S^* \subseteq A_S$ and μ induces a representation μ_{irr} of S_{irr} . It is easy to check that (μ_{irr}, ν) is a $(\text{id} \otimes Q_S)(X)$ -covariant representation.

Therefore, if both A_S and A_T are non-zero, then any X -covariant representation (μ, ν) will induce a X_{irr} -covariant representation.

Theorem 4.9 *Let (T, X, S) and (S, Y, R) be Fourier dualities. Let (μ, ν) and (ρ, λ) be a X -covariant and a Y -covariant representations respectively. Then*

$$e_\mu = e_\lambda \quad \text{as well as} \quad A_S = e_\mu \cdot S^* = \mu^*(\mathcal{L}(H)_*)$$

and hence $A_S \neq (0)$. Moreover, $e_\mu \cdot S^$ is the smallest left Hopf ideal as well as the smallest right Hopf ideal of S^* .*

Proof: The first equality follows from Lemma 4.5, Proposition 4.6 and Remark 4.7(b). Now, $e_\mu = e_\lambda$ implies that $e_\mu \cdot S^*$ is a Hopf ideal of S^* (by Proposition 1.5 and Lemma 4.5) and equals A_S by Lemma 4.8. Moreover, $e_\mu \cdot S^* = e_\lambda \cdot S^*$ is contained in any left or right Hopf ideal of S^* (by Propositions 4.6 and 1.5). This completes the proof.

Corollary 4.10 (a) *With the notations as in Theorem 4.9, μ (respectively, λ) induces a Hopf $*$ -isomorphism from S_{irr} to $\mu(S)$ (respectively, $\lambda(S)$).*

(b) *Let V and W be two C^* -multiplicative unitaries such that $\hat{S}_W = S_V$ (see Example 3.2(a)). Then S_V is irreducible with non-zero Fourier algebra.*

(c) *Let V be any regular irreducible multiplicative unitary and $(\hat{S}_p, W, S_V)$ be the strong dual of S_V . If (μ, ν) is any W -covariant representation, then $\nu((\hat{S}_V)_p) \cong \hat{S}_V$ (as Hopf C^* -algebras). In particular, this applies to $(C^*(G), W_G, C_0(G))$.*

Proof: (a) Note that μ is a Hopf $*$ -homomorphism from S to $\mu(S)$ (Proposition 3.10). Moreover, by Theorem 4.9, $\ker \mu = (\mu^*(\mathcal{L}(H)_*))^\top = (A_S)^\top$. Hence, μ induces a Hopf $*$ -isomorphism from S_{irr} to $\mu(S)$.

(b) This part clearly follows from the fact that $A_{S_V} = L_V^*(\mathcal{L}(H)_*)$ (Theorem 4.9) which separates points of S_V .

(c) Since V is regular, $(\hat{S}_p, W, S_V)$ is a Fourier duality (by Proposition 3.5). Part (c) is now a direct application of parts (a) & (b) (note that if $\hat{q}$ is the quotient from $\hat{S}_p$ to $\hat{S}_V$, then $(\hat{q}, \text{id})$ is also a W -covariant representation).

Definition 4.11 *Suppose that (T, X, S) is a Fourier duality. Then (T, X, S) is said to be a Kac-Fourier duality if both S and T are irreducible and there exist both a X -covariant representation and a X^σ -covariant representation.*

Proposition 4.12 *Let (T, X, S) be a Fourier duality.*

(a) *If (T, X, S) is a Kac-Fourier duality, then it is defined by a multiplicative unitary in the usual way.*

(b) *(T, X, S) is a Kac-Fourier duality if and only if there exist a X^σ -covariant representation as well as a faithful X -covariant representation. It is the case if and only if (T, X, S) and (S, X^σ, T) are defined (in the usual sense) by two multiplicative unitaries V and W respectively.*

Proof: (a) Suppose that (μ, ν) is a X -covariant representation. Let V be the multiplicative unitary $(\nu \otimes \mu)(X)$ (see Proposition 3.10). Then because $\mu^*(\mathcal{L}(H)_*) = A_S$ and $\nu^*(\mathcal{L}(H)_*) = A_T$ (by Theorem 4.9), we have, by Proposition 4.3,

$$\mu(S) = \{\mu((f \otimes \text{id})(X)) : f \in A_T\} = \{(\omega \circ \nu \otimes \mu)(X) : \omega \in \mathcal{L}(H)_*\} = S_V$$

and $\nu(T) = \hat{S}_V$. Moreover, Corollary 4.10(a) implies that both μ and ν are injective. Therefore, μ and ν are Hopf $*$ -isomorphisms from S and T to S_V and $\hat{S}_V$ respectively (see Proposition 3.10).

(b) As above, any X -covariant representation of a Kac-Fourier duality is faithful. Conversely, if there exist a X -covariant representation as well as a X^σ -covariant representation such that one of them is faithful, then Corollary 4.10(a) shows that both S and T are irreducible. Finally, one direction of the second statement clearly follows from part (a) while the other follows from the argument of Corollary 4.10(b).

Note that the multiplicative unitaries obtained above need not be semi-regular nor manageable. Hence Kac-Fourier duality is a far more general framework than Kac system.

Finally, if S is dualizable, we also have the following uniqueness type theorem for the “dual object”.

Theorem 4.13 *Let (T, X, S) , (S, Y, R) , (μ, ν) and (ρ, λ) be the same as in Theorem 4.9. Suppose that S is dualizable. Then $\rho \circ q_R$ induces a Hopf $*$ -isomorphism from $(\hat{S}_p)_{irr}$ to $\rho(R)$ (where q_R is the canonical quotient map from $\hat{S}_p$ to R ; see Proposition 3.5). The same is true for $\nu \circ q_T$. Therefore, if both ρ and ν are injective, then $R \cong T$ (as Hopf C^* -algebras) and Y is the image of X^σ under this isomorphism.*

Proof: Note that $(\mu, \nu \circ q_T)$ is a V_S -covariant representation while $(\rho \circ q_R, \lambda)$ is a V_S^σ -covariant representation (the strong dual $(\hat{S}_p, V_S, S)$ is a Fourier duality because of Proposition 3.5). Corollary 4.10(a) will now give the first two statements. The final statement follows from Proposition 3.5.

The above theorem applies, in particular, to Kac-Fourier dualities. This also gives the following corollary.

Corollary 4.14 *(a) Let V and W be two C^* -multiplicative unitaries (in particular, regular or manageable). If $S_V \cong \hat{S}_W$ as Hopf C^* -algebra, then $\hat{S}_V \cong S_W$ as Hopf C^* -algebras and V is the image of W^σ under these isomorphisms. Moreover, both S_V and $\hat{S}_V$ are irreducible. In this case, if U is another C^* -multiplicative unitary such that $S_V \cong S_U$, then $\hat{S}_V \cong \hat{S}_U$ and U is the image of V under these isomorphisms.*

(b) Let V and W be two semi-regular and balanced (or regular) multiplicative unitaries such that V is irreducible. If $S_V \cong S_W$ as Hopf C^ -algebras, then $\hat{S}_V \cong \hat{S}_W$ as Hopf C^* -algebras and V is the image of W under these isomorphisms.*

References

- [1] S. Baaĵ, Representation reguli re du groupe quantique des d placements de Woronowicz, *Asterisque* (1995), no. 232, 11-48.
- [2] S. Baaĵ and G. Skandalis, Unitaires multiplicatifs et dualit  pour les produits crois s de C^* -alg bres, *Ann. scient.  c. Norm. Sup.*, 4  s rie, t. 26 (1993), 425-488.
- [3] E. Bedos, G.J. Murphy and L. Tuset, Amenability and co-amenable of algebraic quantum groups, to appear in *Int. J. Math. Math. Sci.*, (arXiv:math.OA/0111025).
- [4] Jean De Canni re and Ronny Rousseau, The Fourier Algebra as an Order Ideal of the Fourier-Stieltjes Algebra, *Math. Z.* 186 (1984), 501-507.
- [5] P. Desmedt, J. Quaegebeur and S. Vaes, Amenability and the bicrossed product construction, preprint, (arXiv:math.QA/0111320).
- [6] M. Enock and J.-M. Schwartz, Kac algebras and duality of locally compact groups (Springer-Verlag, 1992).
- [7] V. de M. Iorio, Hopf C^* -algebras and locally compact groups, *Pacific J. Math.* 87 (1980), 75-96.
- [8] J. Kustermans and S. Vaes, Locally compact quantum groups, *Ann. Sci.  cole Norm. Sup.* (4) 33 (2000), no. 6, 837-934.
- [9] J. Kustermans and S. Vaes, Locally compact quantum groups in the von Neumann algebraic setting, *Mathematica Scandinavica*, to appear, (arXiv:math.OA/0005219).

- [10] C. K. Ng, Coactions and crossed products of Hopf C^* -algebras, *Proc. London Math. Soc.*(3) 72(1996), 638-656.
- [11] C. K. Ng, Morphisms of multiplicative unitaries, *Journal of Operator Theory*, 38 (1997), 203–224.
- [12] C. K. Ng, Amenability of Hopf C^* -algebras, *Operator Theoretical Methods (Timișoara, 1998)*, 269-284, Theta Found., Bucharest (2000).
- [13] C. K. Ng, Amenable representations and Reiter's property for Kac algebras, *J. Fun. Anal.* 187 (2001), 163-182.
- [14] C. K. Ng, Examples of Amenable Kac system, *Proc. Amer. Math. Soc.*, to appear, (arXiv: math.OA/0010263).
- [15] M. Takesaki, *Theory of Operator algebras I* (Springer- Verlag, 1979).
- [16] J. -M. Vallin, C^* -algèbres de Hopf et C^* -algèbres de Kac, *Proc. London. Math. Soc.*(3) 50(1985) 131-174.
- [17] Stefaan Vaes and Leonid Vainerman, Extensions of locally compact quantum groups and the bicrossed product construction, preprint, (arXiv:math.OA/0101133).
- [18] Stefaan Vaes and Alfons Van Daele, Hopf C^* -algebras, *Proc. Lond. Math. Soc.* (3) 82, 2001, 337-384.
- [19] S. L. Woronowicz, Unbounded elements affiliated with C^* -algebras and noncompact quantum groups, *Comm. Math. Phys.* 136 (1991), no.2, 399-432.
- [20] S. L. Woronowicz, C^* -algebras generated by unbounded elements, *Reviews in Math. Phys.* 7(1995), no.3, 481-521.
- [21] S. L. Woronowicz, From multiplicative unitaries to quantum groups, *Internat. J. Math.* 7(1996), no.1, 127-149.

Department of Pure Mathematics, Queen's University Belfast, Belfast BT7 1NN, United Kingdom.
 e-mail address: c.k.ng@qub.ac.uk